

Strong one-pion decay of ground state charmed baryons

C. Albertus¹, E. Hernández², J. Nieves¹ and J. M. Verde-Velasco²

¹Departamento de Física Moderna, Universidad de Granada, E-18071 Granada, Spain.

²Grupo de Física Nuclear, Facultad de Ciencias, E-37008 Salamanca, Spain.


Outline of the talk

- Baryon description
 - Three–body Hamiltonian
 - Quark–quark potentials
 - Orbital wave functions
 - Baryon states
- One–pion emission amplitude model
- Results
 - $\Sigma_c \rightarrow \Lambda_c \pi$
 - $\Sigma_c^* \rightarrow \Lambda_c \pi$
 - $\Xi_c^* \rightarrow \Xi_c \pi$
- Summary

Three–body Hamiltonian

$$H = -\frac{\vec{\nabla}_{\vec{R}}^2}{2\overline{M}} + H_{int} \quad H_{int} = \overline{M} + \sum_{j=1,2} H_j^{sp} + V_{l_1, l_2}(\vec{r}_1 - \vec{r}_2, spin) - \frac{\vec{\nabla}_1 \cdot \vec{\nabla}_2}{m_h}$$

$$H_j^{sp} = -\frac{\vec{\nabla}_j^2}{2\mu_j} + V_{h, l_j}(\vec{r}_j, spin), \quad \overline{M} = m_{l_1} + m_{l_2} + m_h, \quad \frac{1}{\mu_j} = \frac{1}{m_{l_j}} + \frac{1}{m_h}$$



Quark–quark Potentials

We use 5 different quark–quark potentials taken from

B. Silvestre-Brac, Few-Body Systems 20, 1 (1996)

R.K. Bhaduri, L.E. Cohler, Y. Nogami, Nuovo Cimento A65, 376 (1981)

Common structure:

- Confinement term
- OGE
 - Coulomb term
 - Hyperfine terms

$$V_{ij}^{q\bar{q}}(r) = -\frac{\kappa(1-e^{-r/r_c})}{r} + \lambda r^p - \Lambda$$
$$+ \left\{ a_0 \frac{\kappa}{m_i m_j} \frac{e^{-r/r_0}}{r r_0^2} + \frac{2\pi}{3m_i m_j} \kappa' (1 - e^{-r/r_c}) \frac{e^{-r^2/x_0^2}}{\pi^{3/2} x_0^3} \right\} \vec{\sigma}_i \vec{\sigma}_j$$
$$V_{ij}^{qq}(r) = \frac{1}{2} V_{ij}^{q\bar{q}}(r)$$

Differences:

- $p = 1$ (lattice QCD) or $p = 2/3$ (Regge trajectories)
- Form factors in OGE terms

Parameters adjusted to light and heavy-light meson spectroscopy

Orbital wave functions

Variational Ansatz $\phi_{l_1, l_2, h}^{S_l}(\vec{r}_1, \vec{r}_2) = N \phi_{l_1}^h(r_1) \phi_{l_2}^h(r_2) F_{l_1, l_2}^{S_l}(r_{12})$

- S_l : Total spin of the light degrees of freedom assumed to be fixed (Heavy Quark Symmetry limit)
- $\phi_{l_j}^h(r_j) = (1 + \alpha_j r_j) \psi_{l_j}^h(r_j)$
Ground state wave function $\psi_{l_j}^h(r_j)$ for the relative motion of the light–quark heavy–quark system corrected at large distances
- Jastrow correlation function

$$F_{l_1, l_2}^{S_l}(r_{12}) = f_{l_1, l_2}^{S_l}(r_{12}) \sum_{j=1}^4 a_j e^{-b_j^2 (r_{12} + d_j)^2} \quad a_1 = 1$$

$$f_{l_1, l_2}^{S_l}(r_{12}) = \begin{cases} 1 - e^{-c r_{12}} & \text{if } V_{l_1, l_2}(r_{12} = 0, S_l) \gg 0 \\ 1 & \text{if } V_{l_1, l_2}(r_{12} = 0, S_l) \leq 0 \end{cases}$$

In momentum space:

$$\tilde{\phi}_{l_1, l_2, h}^{S_l}(\vec{Q}_1, \vec{Q}_2) = \frac{1}{(2\pi)^{3/2}} \int d^3 P N \tilde{\phi}_{l_1}^h(|\vec{P}|) \tilde{\phi}_{l_2}^h(|\vec{Q}_1 + \vec{Q}_2 - \vec{P}|) \tilde{F}_{l_1, l_2}^{S_l}(|\vec{Q}_1 - \vec{P}|)$$

Baryon states

$$\left| B, s \vec{P} \right\rangle_{NR} = \int d^3 Q_1 \int d^3 Q_2 \frac{1}{\sqrt{2}} \sum_{\alpha_1, \alpha_2, \alpha_3} \frac{\hat{\psi}_{\alpha_1, \alpha_2, \alpha_3}^{(B, s)}(\vec{Q}_1, \vec{Q}_2)}{(2\pi)^3 \sqrt{2E_{f_1}(|\vec{p}_1|)2E_{f_2}(|\vec{p}_2|)2E_{f_3}(|\vec{p}_3|)}} \\ \times \left| \alpha_1 \vec{p}_1 = \frac{m_{f_1}}{M} \vec{P} + \vec{Q}_1 \right\rangle \left| \alpha_2 \vec{p}_2 = \frac{m_{f_2}}{M} \vec{P} + \vec{Q}_2 \right\rangle \left| \alpha_3 \vec{p}_3 = \frac{m_{f_3}}{M} \vec{P} - \vec{Q}_1 - \vec{Q}_2 \right\rangle$$

with

$$\alpha_j \equiv (s_j, f_j, c_j), \quad \langle \alpha' \vec{p}' | \alpha \vec{p} \rangle = \delta_{\alpha, \alpha'} (2\pi)^3 2E(|\vec{p}|) \delta^3(\vec{p} - \vec{p}')$$

The normalization is given by

$${}_{NR} \langle B, s' \vec{P}' | B, s \vec{P} \rangle_{NR} = \delta_{s, s'} (2\pi)^3 \delta^3(\vec{P} - \vec{P}')$$

An example:

$$\hat{\psi}_{\alpha_1, \alpha_2, \alpha_3}^{(\Xi_c^0, s)}(\vec{Q}_1, \vec{Q}_2) = \frac{\varepsilon_{c_1 c_2 c_3}}{\sqrt{3!}} (1/2, 1/2, 0; s_1, s_2, 0) \\ \times \frac{\delta_{s_3, s} \delta_{f_3, c}}{\sqrt{2}} \left(\delta_{f_1, d} \delta_{f_2, s} \tilde{\phi}_{d, s, c}^{S_l=0}(\vec{Q}_1, \vec{Q}_2) - \delta_{f_1, s} \delta_{f_2, d} \tilde{\phi}_{s, d, c}^{S_l=0}(\vec{Q}_1, \vec{Q}_2) \right)$$

Pion emission amplitude

PCAC:

$$\partial^\mu J_{A\mu}^{du}(x) = f_\pi m_\pi^2 \Phi_{\pi^-}(x) , \quad f_\pi = 130.7 \text{ MeV}$$

- Matrix elements

$$\left\langle B', s' \vec{P}_{B'} \mid q^\mu J_{A\mu}^{du}(0) \mid B, s \vec{P}_B \right\rangle = -i f_\pi \frac{m_\pi^2}{q^2 - m_\pi^2} \mathcal{A}_{BB'\pi}^{(s,s')}(P_B, P_{B'})$$

$\mathcal{A}_{BB'\pi}^{(s,s')}(P_B, P_{B'})$: Pion emission amplitude

- Pion-pole contribution

$$\left\langle B', s' \vec{P}_{B'} \mid q^\mu J_{A\mu}^{du}(0) \mid B, s \vec{P}_B \right\rangle_{pion-pole} = -i f_\pi \frac{q^2}{q^2 - m_\pi^2} \mathcal{A}_{BB'\pi}^{(s,s')}(P_B, P_{B'})$$

- Non-pole contribution

$$\left\langle B', s' \vec{P}_{B'} \mid q^\mu J_{A\mu}^{du}(0) \mid B, s \vec{P}_B \right\rangle_{non-pole} = i f_\pi \mathcal{A}_{BB'\pi}^{(s,s')}(P_B, P_{B'})$$

$\Sigma_c \rightarrow \Lambda_c \pi$ decay

$$\begin{aligned}
\mathcal{A}_{\Sigma_c^{++} \Lambda_c^+ \pi^+}^{(s,s')}(P, P') &= \frac{-i}{f_\pi} \left\langle \Lambda_c^+, s' \vec{P}' \mid q^\mu J_A^{d,u}(0) \mid \Sigma_c^{++}, s \vec{P} \right\rangle_{non-pole} \\
&= ig_{\Sigma_c^{++} \Lambda_c^+ \pi^+} \bar{u}_{\Lambda_c^+ s'}(\vec{P}') \gamma_5 u_{\Sigma_c^{++} s}(\vec{P})
\end{aligned}$$

$$\Gamma(\Sigma_c^{++} \rightarrow \Lambda_c^+ \pi^+) = \frac{|\vec{q}|}{8\pi M_{\Sigma_c^{++}}^2} g_{\Sigma_c^{++} \Lambda_c^+ \pi^+}^2 \left((M_{\Sigma_c^{++}} - M_{\Lambda_c^+})^2 - m_\pi^2 \right)$$

Using $\vec{P}' = -|\vec{q}|\hat{k}$ and taking into account the different normalization of our NR states

$$\begin{aligned}
g_{\Sigma_c^{++} \Lambda_c^+ \pi^+} &= \frac{-1}{f_\pi} \frac{\sqrt{E_{\Lambda_c^+}(|\vec{q}|) + M_{\Lambda_c^+}} \sqrt{2M_{\Sigma_c^{++}} 2E_{\Lambda_c^+}(|\vec{q}|)}}{|\vec{q}| \sqrt{2M_{\Sigma_c^{++}}}} \\
&\times \left((M_{\Sigma_c^{++}} - E_{\Lambda_c^+}(|\vec{q}|)) A_{\Sigma_c^{++} \Lambda_c^+, 0}^{1/2, 1/2} + |\vec{q}| A_{\Sigma_c^{++} \Lambda_c^+, 3}^{1/2, 1/2} \right)
\end{aligned}$$

$$A_{\Sigma_c^{++} \Lambda_c^+, \mu}^{1/2, 1/2} = {}_{NR} \left\langle \Lambda_c^+, 1/2 - |\vec{q}| \vec{k} \mid J_A^{d,u}(0) \mid \Sigma_c^{++}, 1/2 \vec{0} \right\rangle_{NR, non-pole}$$

$\Sigma_c \rightarrow \Lambda_c \pi$ decay: Results

	$g_{\Sigma_c^{++} \Lambda_c^+ \pi^+}$	$\Gamma(\Sigma_c^{++} \rightarrow \Lambda_c^+ \pi^+)$ [MeV]	$\Gamma(\Sigma_c^+ \rightarrow \Lambda_c^+ \pi^0)$ [MeV]	$\Gamma(\Sigma_c^0 \rightarrow \Lambda_c^+ \pi^-)$ [MeV]
This work	$21.73 \pm 0.32 \pm 0.08$	$2.41 \pm 0.07 \pm 0.02$	$2.79 \pm 0.08 \pm 0.02$	$2.37 \pm 0.07 \pm 0.02$
Experiment		$2.3 \pm 0.2 \pm 0.3^1)$ $2.05^{+0.41}_{-0.38} \pm 0.38^3)$	< 4.6 (CL=90%) ²⁾	$2.5 \pm 0.2 \pm 0.3^1)$ $1.55^{+0.41}_{-0.37} \pm 0.38^3)$
Theory				
CQM		$1.31 \pm 0.04^4)$ $2.025^{+1.1345}_{-0.987}$	$1.31 \pm 0.04^4)$	$1.31 \pm 0.04^4)$ $1.939^{+1.1145}_{-0.954}$
HHCPT	$22, 29.3^6)$	$2.47, 4.38^6)$ $2.5^7)$	$2.85, 5.06^6)$ $3.2^7)$	$2.45, 4.35^6)$ $2.4^7)$ $1.94 \pm 0.57^8)$
LFQM		$1.64^9)$	$1.70^9)$	$1.57^9)$
RTQM		$2.85 \pm 0.19^{10})$	$3.63 \pm 0.27^{10})$	$2.65 \pm 0.19^{10})$

1) M. Artuso *et al.* (CLEO Coll.), Phys. Rev. D65,071101 (2002)

2) R. Ammar *et al.* (CLEO Coll.), Phys. Rev. Lett. 86, 1167 (2001)

3) J. M. Link *et al.* (FOCUS Coll.), Phys. Lett. B525, 205 (2002)

4) J. L. Rosner, Phys. Rev. D52, 6461 (1995)

5) D. Pirjol, T. M. Yan, Phys. Rev. D56, 5483 (1997)

6) T. M. Yan *et al.*, Phys. Rev. D46, 1148 (1992)

7) M.-Q. Huang, Y.-B. Dai, C.-S. Huang, Phys. Rev. D52, 2986 (1995)

8) H.-Y. Cheng, Phys. Lett. B399, 281 (1997)

9) S. Tawfig, P. J. O'Donnell, J. G. Körner, Phys. Rev. D56, 054010 (1998)

10) M. A. Ivanov, J. G. Körner, V. E. Lyubovitskij, A. G. Rusetsky, Phys. Rev. D46, 1148 (1999); Phys. Lett. B442, 435 (1998)

Σ_c^* → $\Lambda_c \pi$ decay

$$\begin{aligned}
\mathcal{A}_{\Sigma_c^{*++} \Lambda_c^+ \pi^+}^{(s,s')}(P, P') &= \frac{-i}{f_\pi} \left\langle \Lambda_c^+, s' \vec{P}' \mid q^\mu J_A^d u_\mu(0) \mid \Sigma_c^{*++}, s \vec{P} \right\rangle_{non-pole} \\
&= i \frac{g_{\Sigma_c^{*++} \Lambda_c^+ \pi^+}}{2M_{\Lambda_c^+}} q_\nu \bar{u}_{\Lambda_c^+ s'}(\vec{P}') u_{\Sigma_c^{*++} s}^\nu(\vec{P})
\end{aligned}$$

$$\Gamma(\Sigma_c^{*++} \rightarrow \Lambda_c^+ \pi^+) = \frac{|\vec{q}|^3}{24\pi M_{\Sigma_c^{*++}}^2} \frac{g_{\Sigma_c^{*++} \Lambda_c^+ \pi^+}^2}{4M_{\Lambda_c^+}^2} \left((M_{\Sigma_c^{*++}} + M_{\Lambda_c^+})^2 - m_\pi^2 \right)$$

$$\begin{aligned}
g_{\Sigma_c^{*++} \Lambda_c^+ \pi^+} &= \frac{\sqrt{3}}{f_\pi \sqrt{2}} \frac{2M_{\Lambda_c^+} \sqrt{2M_{\Sigma_c^{*++}} 2E_{\Lambda_c^+}(|\vec{q}|)}}{|\vec{q}| \sqrt{2M_{\Sigma_c^{*++}} (E_{\Lambda_c^+}(|\vec{q}|) + M_{\Lambda_c^+})}} \\
&\times \left((M_{\Sigma_c^{*++}} - E_{\Lambda_c^+}(|\vec{q}|)) A_{\Sigma_c^{*++} \Lambda_c^+, 0}^{1/2, 1/2} + |\vec{q}| A_{\Sigma_c^{*++} \Lambda_c^+, 3}^{1/2, 1/2} \right)
\end{aligned}$$

$$A_{\Sigma_c^{*++} \Lambda_c^+, \mu}^{1/2, 1/2} = {}_{NR} \left\langle \Lambda_c^+, 1/2 - |\vec{q}| \vec{k} \mid J_A^d u_\mu(0) \mid \Sigma_c^{*++}, 1/2 \vec{0} \right\rangle_{NR, non-pole}$$

$\Sigma_c^* \rightarrow \Lambda_c \pi$ decay: Results

	$g_{\Sigma_c^* ++ \Lambda_c^+ \pi^+}$	$\Gamma(\Sigma_c^* ++ \rightarrow \Lambda_c^+ \pi^+)$ [MeV]	$\Gamma(\Sigma_c^* + \rightarrow \Lambda_c^+ \pi^0)$ [MeV]	$\Gamma(\Sigma_c^* 0 \rightarrow \Lambda_c^+ \pi^-)$ [MeV]
This work	$36.20 \pm 0.75 \pm 0.13$	$17.52 \pm 0.74 \pm 0.12$	$17.31 \pm 0.73 \pm 0.12$	$16.90 \pm 0.71 \pm 0.12$
Experiment		$14.1^{+1.6}_{-1.5} \pm 1.4^{11})$	< 17 (CL=90%) ²⁾	$16.6^{+1.9}_{-1.7} \pm 1.4^{11})$
Theory				
QCDSR	$13.8 \div 24.2^{12})$			
	$32.5 \pm 2.1 \pm 6.9^{13})$			
CQM		$20^{4)}$	$20^{4)}$	$20^{4)}$
HHCPT		$25^{7})$	$25^{7})$	$25^{7})$
LFQM		$12.84^{9})$		$12.40^{9})$
RTQM		$21.99 \pm 0.87^{10})$		$21.21 \pm 0.81^{10})$

11) S. B. Athar *et al.* (CLEO Coll.), Phys. Rev. D71, 051101 (2005)

12) A. G. Grozin, O. I. Yakovlev, Eur. Phys. J. C2, 721 (1998)

13) S.-L. Zhu, Y.-B. Dai, Phys. Lett. B429, 72 (1998)

Ξ_c^* → $\Xi_c \pi$ decay

$$\begin{aligned} \mathcal{A}_{\Xi_c^* + \Xi_c^0 \pi^+}^{(s,s')}(P, P') &= \frac{-i}{f_\pi} \left\langle \Xi_c^0, s' \vec{P}' | q^\mu J_A^{d,u}(0) | \Xi_c^* +, s \vec{P} \right\rangle_{non-pole} \\ &= i \frac{g_{\Xi_c^* + \Xi_c^+ \pi^+}}{M_{\Xi_c^+} + M_{\Xi_c^0}} q_\nu \bar{u}_{\Xi_c^0 s'}(\vec{P}') u_{\Xi_c^* + s}^\nu(\vec{P}) \end{aligned}$$

$$\Gamma(\Xi_c^* + \rightarrow \Xi_c^0 \pi^+) = \frac{|\vec{q}|^3}{24\pi M_{\Xi_c^* +}^2} \frac{g_{\Xi_c^* + \Xi_c^0 \pi^+}^2}{(M_{\Xi_c^+} + M_{\Xi_c^0})^2} \left((M_{\Xi_c^* +} + M_{\Xi_c^0})^2 - m_\pi^2 \right)$$

$$\begin{aligned} g_{\Xi_c^* + \Xi_c^0 \pi^+} &= \frac{\sqrt{3}}{f_\pi \sqrt{2}} \frac{(M_{\Xi_c^+} + M_{\Xi_c^0}) \sqrt{2M_{\Xi_c^* +} 2E_{\Xi_c^0}(|\vec{q}|)}}{| \vec{q} | \sqrt{2M_{\Xi_c^* +} (E_{\Xi_c^0}(|\vec{q}|) + M_{\Xi_c^0})}} \\ &\times \left((M_{\Xi_c^* +} - E_{\Xi_c^0}(|\vec{q}|)) A_{\Xi_c^* + \Xi_c^0, 0}^{1/2, 1/2} + |\vec{q}| A_{\Xi_c^* + \Xi_c^0, 3}^{1/2, 1/2} \right) \end{aligned}$$

$$A_{\Xi_c^* + \Xi_c^0, \mu}^{1/2, 1/2} = {}_{NR} \left\langle \Xi_c^0, 1/2 - |\vec{q}| \vec{k} | J_A^{d,u}(0) | \Xi_c^* +, 1/2 \vec{0} \right\rangle_{NR, non-pole}$$

Ξ_c^* → $\Xi_c \pi$ decay: Results

	$g_{\Xi_c^* + \Xi_c^0 \pi^+}$	$\Gamma(\Xi_c^* + \rightarrow \Xi_c^0 \pi^+)$ [MeV]	$\Gamma(\Xi_c^* + \rightarrow \Xi_c^+ \pi^0)$ [MeV]	$\Gamma(\Xi_c^{*0} \rightarrow \Xi_c^+ \pi^-)$ [MeV]	$\Gamma(\Xi_c^{*0} \rightarrow \Xi_c^0 \pi^0)$ [MeV]
This work	$-28.83 \pm 0.50 \pm 0.10$	$1.84 \pm 0.06 \pm 0.01$	$1.34 \pm 0.04 \pm 0.01$	$2.07 \pm 0.07 \pm 0.01$	$0.956 \pm 0.030 \pm 0.007$
Theory		$1.12^{9)}$	$0.69^{9)}$	$1.16^{9)}$	$0.72^{9)}$
LFQM					
RTQM		$1.78 \pm 0.33^{10})$	$1.26 \pm 0.17^{10})$	$2.11 \pm 0.29^{10})$	$1.01 \pm 0.15^{10})$
		$\Gamma(\Xi_c^* + \rightarrow \Xi_c^0 \pi^+ + \Xi_c^+ \pi^0)$ [MeV]	$\Gamma(\Xi_c^* + \rightarrow \Xi_c^0 \pi^+ + \Xi_c^+ \pi^0)$ [MeV]		
This work		$3.18 \pm 0.10 \pm 0.01$		$3.03 \pm 0.10 \pm 0.01$	
Exp.		< 3.1 (CL=90%) ¹⁴⁾		< 5.5 (CL=90%) ¹⁵⁾	
Theory					
CQM		$< 2.3 \pm 0.1^{4)}$		$< 2.3 \pm 0.1^{4)}$	
HHCPT		$1.191 \div 3.971^{5})$		$1.230 \div 4.074^{5})$	
LFQM		$2.44 \pm 0.85^{8})$		$2.51 \pm 0.88^{8})$	
RTQM		$1.81^{9})$		$1.88^{9})$	
		$3.04 \pm 0.50^{10})$		$3.12 \pm 0.33^{10})$	

14) L. Gibbons *et al.* (CLEO Coll.), Phys. Rev. Lett. 77, 810 (1996)

15) P. Avery *et al.* (CLEO Coll.), Phys. Rev. Lett. 75, 4364 (1995)

Summary

- First dynamical calculation of these observables within a non relativistic quark model
- Wave functions
 - Obtained using a variational ansatz made possible by HQS constraints
- One-pion emision amplitude
 - We use PCAC to relate the one-pion emission amplitude to weak current matrix elements
- Results
 - Stable against different quark-quark interaction
 - Overall good agreement with experiment